

## APPENDIX 1: Drugs collected for Wastage Study 1.

Amiodarone	Levetiracetam	Potassium chloride
Amphotericin B, liposomal	Lidocaine hydrochloride 1%	Ranitidine
L-Arginine	Magnesium sulphate	Ropivacaine hydrochloride
Caffeine	Methylprednisolone	Sodium bicarbonate
Cefotaxime	Metronidazole	Sodium phosphate
Ceftriaxone	Milrinone lactate	Tobramycin
Cefuroxime	Naloxone	Tuberculin purified protein derivative
Enoxaparin	Novolin ge Toronto	Vancomycin
Fosphenytoin, phenytoin sodium	Penicillin G	Zoledronic acid

## APPENDIX 2: Drugs collected for Wastage Study 2.

Acetylcysteine	Fluconazole	Norepinephrine
Amphotericin B, liposomal	Gentamicin	Pantoprazole
Caffeine	Heparin	Penicillin G
Cefazolin	Levofloxacin	Pentamidine
Cefotaxime	Lidocaine hydrochloride 1%	Piperacillin-tazobactam
Ceftriaxone	Magnesium sulphate	Potassium acetate
Ciprofloxacin	Methylprednisolone	Sodium bicarbonate
Cloxacillin	Metronidazole	Vancomycin
Enoxaparin	Micro Plus 6 Pediatric	Zoledronic acid

## APPENDIX 3: Drugs collected for Wastage Study 3.

Amphotericin B, liposomal	Cefuroxime	Penicillin G
Azithromycin	Ciprofloxacin	Piperacillin
Caspofungin	Clindamycin	Piperacillin-tazobactam
Cefazolin	Enoxaparin	Ranitidine
Cefotaxime	Filgrastim	Tobramycin
Cefoxitin	Fluconazole	Vancomycin
Ceftazidime	Metronidazole	
Ceftriaxone	Pantoprazole	

Supplementary material for Hollis K, Payen F, Vaillancourt R. Implementation of beyond-use date guidelines for single-use vials at a pediatric hospital. *Can J Hosp Pharm.* 2021;74(1):70-4.