

Heparin-Induced Priapism

The authors of a case report of heparin-induced priapism, published in the December issue, state that “the treatment of priapism remains largely surgical, but diphenhydramine, trazodone, cyproheptadine, and benztropine have been used with various degrees of success.”¹

In fact, trazodone is not used as a treatment of priapism, but rather is known to induce priapism.² I believe that the authors have misquoted the reference cited for this information, which states that “diphenhydramine has been employed successfully in priapism secondary to antipsychotics (eg. trazodone and chlorpromazine), and α -adrenergic blockers, and cyproheptadine and benztropine have also been shown to be useful in priapism secondary to trazodone.”²

Suzanne Singh

Class of 2000
Faculty of Pharmacy
University of Toronto
Toronto, Ontario

References

1. Arustei L, Kwan A. Heparin-induced priapism. *Can J Hosp Pharm* 1999;52:378-9.
2. Banos JE, Bosch F, Farre M. Drug-induced priapism: its aetiology, incidence and treatment. *Med Toxicol* 1989;4:46-58.

Reply:

We have reviewed the reference, and Ms Singh is correct. It appears that during the development of the paper an error was introduced into the statement in question. The text should have read as follows:

The treatment of priapism remains largely surgical, but α -adrenoceptor agonists (for example, intracavernosal metaraminol) have been used. Diphenhydramine has also been used successfully in priapism secondary to chlorpromazine, α -adrenoceptor blockers and trazodone, and cyproheptadine and benztropine have been

employed successfully in trazodone-induced priapism. None of these studies was controlled. Further information is available in the review by Banos and colleagues.¹

We regret the error.

Liviu Arustei, BScPhm
Archie Kwan, BSc(Pharm)
Pharmacy Department
Centenary Health Centre
Scarborough, Ontario

Reference

1. Banos JE, Bosch F, Farre M. Drug-induced priapism: its aetiology, incidence and treatment. *Med Toxicol* 1989;4:46-58.

Change in Formulations for Acetazolamide and Allopurinol Suspensions

At The Hospital for Sick Children we have recently modified the way in which we prepare suspensions for acetazolamide and allopurinol. We now make them with the Ora-Plus® and Ora-Sweet® vehicles.

The revised formulas are presented here.

Acetazolamide 25 mg/mL Oral Suspension

Ingredients	Manufacturer	Quantity
Acetazolamide, 250-mg tablets	Apotex*	10
Ora-Plus®	Paddock	50 mL
Ora-Sweet®	Paddock	50 mL
Ora-Plus® – Ora-Sweet® combination	The Hospital for Sick Children	qs 100 mL

* No substitution.

